



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## CHARACTERISING PET DEGRADING IDEONELLA SAKAIENSIS & ENGINEERING OF PETase THROUGH MUTAGENESIS

Jenkins, Samantha Gini; Fonseca, César; Cannella, David; Varrone, Cristiano

DOI (link to publication from Publisher):  
[10.13140/RG.2.2.14822.47684](https://doi.org/10.13140/RG.2.2.14822.47684)

Creative Commons License  
Unspecified

Publication date:  
2019

Document Version  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*  
Jenkins, S. G., Fonseca, C., Cannella, D., & Varrone, C. (2019). *CHARACTERISING PET DEGRADING IDEONELLA SAKAIENSIS & ENGINEERING OF PETase THROUGH MUTAGENESIS*. Poster presented at 5th Edition of The International Conferences Green Chemistry - White Biotechnology on (BIO-)Polymers and Ecocircularity: From Challenges to Opportunities; Bruxelles. 8-9 May 2019..  
<https://doi.org/10.13140/RG.2.2.14822.47684>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/333394674>

# CHARACTERISING PET DEGRADING IDEONELLA SAKAIENSIS & ENGINEERING OF PETase THROUGH MUTAGENESIS

Poster · May 2019

DOI: 10.13140/RG.2.2.14822.47684

CITATIONS

0

READS

2,041

4 authors:



**Samantha G.R. Jenkins**

Aalborg University

3 PUBLICATIONS 0 CITATIONS

[SEE PROFILE](#)



**César Fonseca**

Chr. Hansen A/S

40 PUBLICATIONS 2,327 CITATIONS

[SEE PROFILE](#)



**David Cannella**

Université Libre de Bruxelles

28 PUBLICATIONS 920 CITATIONS

[SEE PROFILE](#)



**Cristiano Varrone**

Aalborg University

169 PUBLICATIONS 854 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Biological depolymerization of plastics [View project](#)



GRAIL Project (EU FP7 KBBE) [View project](#)



# CHARACTERISING PET DEGRADING IDEONELLA SAKAIENSIS & ENGINEERING OF PETase THROUGH MUTAGENESIS



Samantha Jenkins,   Cesar Simones da Fonseca,  David Cannella,  Cristiano Varrone, 

Universite Libre de Bruxelles and Aalborg University

Email: Sjenki17@student.aau.dk, csf@bio.aau.dk, David.cannella@ulb.ac.be, Cva@bio.aau.dk.



## INTRODUCTION

8300 million tonnes of plastic produced since early 20th century. Plastic pollution leads to toxic substances being released into our oceans, land and even air. Methods such as recycling and depolymerisation whilst excellent will only ever result in more plastic.

- 33% used once, and thrown away. Only 8% recycled (Wang et al., 2015).
- 8 million tonnes p/y released into oceans


Degrade into microplastics

- Enter the food chain and are on the rise
- Unexamined health impacts

## PET (POLYETHYLENE TEREPHTHALATE)



- Crystalline plastic
- Drinks bottles, fleece, carpets
- Low melting temp
- Easy reformation
- Only 7% recycled into new products
- Made from ethylene monomers and terephthalic acid



## BIODEGRADATION

- Plastic degrading microorganisms exist in abundance (see table), the process is just too slow
- Biodegradation removes plastic from cycle and can turn plastic waste into a resource
- and result in profitable products (product engineering)

Plastic	Microorganism	Enzyme	Source	Reference
Polyethylene	<i>Brevibacillus borstelensis</i>		Bacterial	[35] [14]
	<i>Rhodococcus ruber</i>		Bacterial	[36] [37] [14]
	<i>Penicillium simplicissimum</i> JK		Bacterial	[38] [14]
	<i>Ideonella sakaiensis</i>	lipase	Bacterial	[6]
Polyethylene adipate (PEA)	<i>Pseudomonas putida</i>	peroxidase	Bacterial	[6]
	<i>Penicillium, Rhizopus arrizus</i>	lipase	Bacterial	[6]
Polyhydroxyalkanoate (PHA)	<i>Pseudomonas stutzeri</i>	Serine hydrolase	Bacterial	[6]
Polyethylene terephthalate	<i>Ideonella sakaiensis</i>	Hydrolase	Bacterial	[6]
Polyurethane	<i>penicillium microspora</i>	Hydrolase	Fungal	[39]
	<i>Comamonas acidovorans</i> TB-33	esterase	Bacterial	[40] [41] [14]
	<i>Penicillium microspora</i>	Hydrolase	Fungal	[39]
	<i>Phanerochaete chrysosporium</i>	Manganese peroxidase	Fungal	
Polyvinyl chloride	<i>Pseudomonas chlororaphis</i>		Bacterial	[14]
	<i>Curvularia reniformis</i>		Fungal	[42] [14]
	<i>Fusarium solani</i>		Fungal	[14]
	<i>Candida glabrata</i> sp.		Fungal	[14]
	<i>Pseudomonas putida</i> AJ		Bacterial	[14]
	<i>Chlorobacterium</i> TD		Bacterial	[14]
Polycaprolactone (PCL)	<i>Pseudomonas fluorescens</i> B-22		Bacterial	[43] [14]
	<i>Aspergillus niger</i>		Bacterial	[14]
	<i>Aspergillus fumigatus</i>	Glycosidase	Fungal	[6]
	<i>Aspergillus niger</i>		Fungal	[14]
	<i>Fusarium</i>		Fungal	[14]
Polybutylene succinate (PBS)	<i>Rhizopus delemar</i>	Lipase	Bacterial	[6]
	<i>Penicillium, Rhizopus arrizus</i>			
	<i>Aspergillus oryzae</i>	cutinase	Fungal	[6]
	<i>Penicillium, Rhizopus arrizus</i>	Lipase	Bacterial	[6]

## IDEONELLA SAKAIENSIS

Most effective plastic (PET) degrading organism. (Oda group, Kyoto Institute of Technology, 2016)

Gram negative, aerobic, rod-shaped, non-spore forming, single flagellated



Image of Ideonella sakaiensis producing tendrils which attach to substrate. (Yoshida et al., 2016)

## PROJECT AIM

Characterise the bacterial strain *Ideonella sakaiensis* for the engineering of its plastic degrading PETase enzyme to increase its catalytic activity using genetic engineering techniques such as directed mutagenesis.

## MATERIALS AND METHODS

### CHARACTERISING I. SAKAIENSIS


Preference to grow on solid medium with PET or PE plastic as attachment of tendrils to substrate is require to begin degradation:




Examples of I.Sakaiensis Colonies; round Clear and small

## ENZYME

- Serine hydrolase, charge relay system
- Coupled with MHETase which breakdown the products further in ethylene glycol and terephthalic acid
- PETase bottleneck of enzyme cascade



Degradation of PET into Ethylene glycol and terephthalic acid by PETase



3D structure of PETase enzyme, (Han et al., 2017)


### DIRECTED MUTAGENESIS

Modifications to enzyme active site improve binding of substrate:


**S238F & W159H mutation:**

- Serine to phenalalynine (S238F) = more aromatic interactions & stable 'docking' for PET
- Wild-type PET stabilised by W185 and W159 interactions only.

Double mutant provides four aromatic interactions to stabilise the PET




Degradation of PET film by WT I. sakaiensis (left) and double mutant (far right) (Austin et al., 2018)



### Arg280 mutation:

- Residue in the binding site (Arg280) = polar and protruding region
- Substituting Arg280 for Alanine, = longer, unhindered, substrate binding.



(Joo et al., 2018)

## FUTURE PERSPECTIVES

- Faster and more broad spectrum plastic biodegradation
- Make plastic into a resource by engineering products
- Bio-sensors: Ocean microplastic problem
- Demobilization onto filters
- Demobilization onto underwater structures (artificial reefs)
- Synthetic consortia

## References:

Austin, H.P., Allen, M.D., Donohoe, B.S., Rorrer, N.A., Kearns, F.L., Silveira, R.L., Pollard, B.C., Dominick, G., Duman, R., El Omari, K. and Mykhaylyk, V., 2018. Characterization and engineering of a plastic-degrading aromatic polyesterase. *Proceedings of the National Academy of Sciences*, 115(19), pp.E4350-E4357

Champakalakshmi Ravi Kyaw Myint Bhone, Sing Lim Chu Sakharkar Meena Kishore, and Sakharkar Kishore R. Biodegradation of Low Density Polythene (LDPE) by Pseudomona Species. *Rivista degli infortuni e delle malattie professionali*, 1951

Han, X., Liu, W., Huang, J.W., Ma, J., Zheng, Y., Ko, T.P., Xu, L., Cheng, Y.S., Chen, C.C. and Guo, R.T., 2017. Structural insight into catalytic mechanism of PET hydrolase. *Nature communications*, 8(1), p.2106.

Joo, S., Cho, I.J., Seo, H., Son, H.F., Sagong, H.Y., Shin, T.J., Choi, S.Y., Lee, S.Y. and Kim, K.J., 2018. Structural insight into molecular mechanism of poly (ethylene terephthalate) degradation. *Nature communications*, 9(1), p.382.

Liu, B., He, L., Wang, L., Li, T., Li, C., Liu, H., Luo, Y. and Bao, R., 2018. Protein Crystallography and Site-Direct Mutagenesis Analysis of the Poly (ethylene terephthalate) Hydrolase PETase from Ideonella sakaiensis. *ChemBioChem*, 19(14), pp.1471-1475.

Obranić, S., Babić, F. and Maravić-Vlahović, G., 2013. Improvement of pBBR1MCS plasmids, a very useful series of broad-host-range cloning vectors. *Plasmid*, 70(2), pp.263-267.

Tanasupawat, S., Takehana, T., Yoshida, S., Hiraga, K. and Oda, K., 2016. Ideonella sakaiensis sp. nov., isolated from a microbial consortium that degrades poly (ethylene terephthalate). *International journal of systematic and evolutionary microbiology*, 66(8), pp.2813-2818.

Yoshida, S., Hiraga, K., Takehana, T., Taniguchi, I., Yamaji, H., Maeda, Y., Toyohara, K., Miyamoto, K., Kimura, Y. and Oda, K., 2016. A bacterium that degrades and assimilates poly (ethylene terephthalate). *Science*, 351(6278), pp.1196-1199.

Yutaka Tokiwa, Buenaventurada Calabia, Charles Ugwu, Seiichi Aiba, Yutaka Tokiwa, Buenaventurada P. Calabia, Charles U. Ugwu, and Seiichi Aiba. Biodegradability of Plastics. *International Journal of Molecular Sciences*, 10(9):3722–3742, 8 2009